

# PHYSICIAN'S ORDERS



Drug Allergies

NAME

ROOM NO.  
(ADDRESS)

HOSP. NO.

PHYSICIAN

## Pre-Operative Orders

Dr. Thomas VanBruggen

1. **Admit to:**

- Outpatient Status
- Inpatient Status

2. **Procedure:** \_\_\_\_\_

3. **Vital Signs:**

- Routine
- Other: \_\_\_\_\_

4. **Activity:**

- Up ad lib
- Bedrest
- Bedrest with BRPs
- Up with assistance

5. **Diet:**

- NPO
- Clear liquids
- Other: \_\_\_\_\_
- NPO after 2400

6. **I&O:**  Yes  No

7. **IV Lactated Ringers** @ \_\_\_\_\_ cc/hr

8. **Foley Catheter:**

- Yes
- Place in OR

9. **RT Consult:**

- Yes
- Smoking Cessation

10. **Beta Blocker Therapy:**

- Not on beta block pre-operatively
- Patient on beta blocker. Administer usual morning dose with sip of water: \_\_\_\_\_ mg @ \_\_\_\_\_
- Patient on a beta blocker on admission but not ordered for perioperative period. Contraindications of: \_\_\_\_\_

11. **DVT prophylaxis:**

- Lovenox 40 mg subcutaneous pre-operatively
- Have pharmacy calculate Lovenox dosing

*In addition to or may be used exclusively if documented reason for not using pharmacological prophylaxis:*

- Graduated compression socks
- Intermittent pneumatic devices

*Reason for no pharmacological prophylaxis:*

\_\_\_\_\_

12. **Antibiotic prophylaxis:**

*Lower gastrointestinal procedures:*

- Cefoxitin 2 grams IV per anesthesia pre-operatively
- Neomycin, E-mycin base 1 gm each @ 1400, 1500 & 2300 day prior to surgery with Cefoxitin 2 gm IV per anesthesia pre-operatively

If  $\beta$ -lactam allergy:

- Clindamycin 900 mg IV per anesthesia pre-operatively

*Upper gastrointestinal, gynecologic, head and neck procedures:*

- Cefazolin pre-operatively

\*If patient less than 80 kg, 1 gm IV per anesthesia

\*If patient 80 Kg or greater, 2 gm IV per anesthesia

If  $\beta$ -lactam allergy:

- Clindamycin 900 mg IV per anesthesia pre-operatively

*Hip, orthopedic, vascular*

- Cefazolin pre-operatively

\*If patient less than 80 kg, 1 gm IV per anesthesia

\*If patient 80 Kg or greater, 2 gm IV per anesthesia

If  $\beta$ -lactam allergy:

- Clindamycin 900 mg IV per anesthesia pre-operatively

Other: \_\_\_\_\_

- Vancomycin - Pharmacy to calculate dose and levels

If Vancomycin, must have documented indication for use:

\_\_\_\_\_

13. **Other:** \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

PHYSICIAN SIGNATURE

DATE/TIME